



ACC.14

TCT@ACC-12 | innovation in intervention

A1307

JACC April 1, 2014

Volume 63, Issue 12



## Prevention

### DIFFERENTIAL IMPACT OF RENAL SYMPATHETIC DENERVATION ON INDEXES OF SHORT-TERM BLOOD PRESSURE VARIABILITY IN PATIENTS WITH RESISTANT HYPERTENSION

Poster Contributions

Hall C

Saturday, March 29, 2014, 10:00 a.m.-10:45 a.m.

Session Title: New Developments in Renal Denervation

Abstract Category: 21. Prevention: Hypertension

Presentation Number: 1111-132

Authors: *Konstantinos P. Tsioufis, Vasilios Papademetriou, Matthew Worthley, Derek Chew, Ajay Sinhal, Ian Meredith, Yuvaraj Malaipapan, Stephen G. Worthley, First Cardiology Clinic, University of Athens, Hippokration Hospital, Athens, Greece, Cardiovascular Res Ctr, Royal Adelaide Hosp and Dept of Med, Univ of Adelaide, Adelaide, Australia*

**Background:** Transluminal renal sympathetic denervation (RDN) reduces blood pressure (BP) in patients with drug resistant hypertension (RHTN). We assessed the effect of RDN on indexes of short-term blood pressure variability (BPV) in patients with drug RHTN.

**Methods:** In the EnligHTN I study, patients with drug-RHTN underwent office and ambulatory BP measurements (ABPM) at baseline and 6 months after RDN using the multi-electrode EnligHTN ablation catheter (St. Jude Medical, USA). We analyzed BP data for 31 patients (office BP 178.3/94.3 mmHg, 24-hour BP 147.5/81.2 mmHg). For each patient the standard deviation (SD) was recorded of all systolic and diastolic BP during 24-hour, daytime and nighttime and weighted 24-hour BP SD. Average real variability (ARV) of 24-hour systolic and diastolic BP and the time rate of systolic and diastolic BP variation defined as the first derivative of the BP values against time.

**Results:** There was a significant reduction in 24-hour BP by 10.2/6 mmHg, ( $p < 0.001$  for both SBP and DBP) and in maximum values of systolic and diastolic BP from 185/106.5 to 170.8/97.7 mmHg ( $p < 0.001/0.001$  respectively) 6 months post RDN. There were no significant changes in SDs of 24-hour systolic and diastolic BP (from 15.9/10.5 to 15.9/10.8 mmHg), as well as in the daytime and nighttime SDs ( $p = \text{NS}$  for all). There was no difference in the weighted SD of 24-hour systolic and diastolic BP (from 13.7/9.2 to 13.5/9.2 mmHg), which adjusts the SD for the nocturnal BP fall. Additionally, ARV of 24-hour systolic and diastolic BP remained practically unchanged (from 10.1/7.4 to 10.3/7.5 mmHg). In contrast, the rates of systolic and diastolic 24-hour BP variation were significantly decreased (from 0.40/0.30 to 0.34/0.24,  $p = 0.030/0.006$ ). Similarly, a significant variation was observed in the daytime and nighttime rates of systolic and diastolic variation after RDN.

**Conclusion:** RDN exerts a differential impact on short-term BPV indexes, as assessed by ABPM, in patients with drug RHTN. Specifically, while conventional BPV indexes seem to be practically unaffected after RDN, a significant decrease in the rate of BP variation is observed 6 months after RDN.